

APPENDIX 1

EAST KENT

Indicator	Compared to England
Life expectancy at birth (Male)	Similar
Life expectancy at birth (Female)	Similar
Smoking Prevalence in adults (18+) - current smokers (APS)	Similar
Percentage of adults (aged 18+) classified as overweight or obese	Similar
Children with excess weight Year 6, three year average	Similar
Percentage of physically inactive adults	Better
Admission episodes for alcohol-specific conditions	Better
Air pollution: fine particulate matter	Not compared
Total number of prescribed antibiotic items per STAR-PU	Higher
Breast cancer screening coverage (females aged 50-70)	Better
Cervical cancer screening coverage (females aged 25-49)	Better
Bowel cancer screening coverage (persons aged 60-74)	Better
Infant mortality rate	Similar
Low birth weight of term babies	Better
Stillbirth rate	Similar
Smoking status at time of delivery	Worse
AE attendances (0-4 years)	Worse
Percentage of 5 year olds with experience of visually obvious dental decay	Better

WEST KENT

Indicator	Compared to England
Life expectancy at birth (Male)	Better
Life expectancy at birth (Female)	Better
Smoking Prevalence in adults (18+) - current smokers (APS)	Better
Percentage of adults (aged 18+) classified as overweight or obese	Similar
Children with excess weight Year 6, three year average	Better
Percentage of physically inactive adults	Better
Admission episodes for alcohol-specific conditions	Better
Air pollution: fine particulate matter	Not compared
Total number of prescribed antibiotic items per STAR-PU	Higher
Breast cancer screening coverage (females aged 50-70)	Better
Cervical cancer screening coverage (females aged 25-49)	Better
Bowel cancer screening coverage (persons aged 60-74)	Better
Infant mortality rate	Better
Low birth weight of term babies	Better
Stillbirth rate	Similar
Smoking status at time of delivery	Worse
AE attendances (0-4 years)	Worse
Percentage of 5 year olds with experience of visually obvious dental decay	Better

Indicator	Compared to England
Under 18s conception rate / 1,000	Better
Emergency hospital admissions for asthma (< 19 yrs)	Better
Emergency hospital admissions for epilepsy (< 19 yrs)	Better
Emergency hospital admissions for diabetes (< 19 yrs)	Similar
Hospital admissions for mental health conditions (0-17 years)	Better
Hospital admissions as a result of self-harm (10-24 years)	Worse
Hospital admissions due to substance misuse (15-24 years)	Similar
Hypertension: QOF prevalence (all ages)	Lower
Diabetes: QOF prevalence (17+)	Lower
CHD: QOF prevalence (all ages)	Lower
CKD: QOF prevalence (18+)	Higher
Stroke: QOF prevalence (all ages)	Similar
Deaths from circulatory disease, under 75 years	Better
Deaths from all cancer, under 75 years	Better
Cancer diagnosed at early stage (experimental statistics)	Not compared
Unplanned hospitalisation for chronic ACSC	Better
Depression: Recorded prevalence (aged 18+)	Higher
Serious Mental Illness: QOF prevalence (all ages)	Lower

Indicator	Compared to England
Suicide rate (Persons)	Similar
Suicide rate (Male)	Similar
Estimated dementia diagnosis rate (aged 65 and over)	Not compared
Emergency hospital admissions due to falls (persons aged 65 and over)	Worse
Emergency hospital admissions for hip fracture (persons aged 65 and over)	Similar
Osteoporosis: QOF prevalence (50+)	Higher

DARTFORD GRAVESHAM AND SWANLEY

Indicator	Compared to England
Life expectancy at birth (Male)	Similar
Life expectancy at birth (Female)	Similar
Smoking Prevalence in adults (18+) - current smokers (APS)	Worse
Percentage of adults (aged 18+) classified as overweight or obese	Worse
Children with excess weight Year 6, three year average	Worse
Percentage of physically inactive adults	Worse
Admission episodes for alcohol-specific conditions	Better
Air pollution: fine particulate matter	Not compared
Total number of prescribed antibiotic items per STAR-PU	Similar
Breast cancer screening coverage (females aged 50-70)	Worse
Cervical cancer screening coverage (females aged 25-49)	Similar
Bowel cancer screening coverage (persons aged 60-74)	Similar
Infant mortality rate	Similar
Low birth weight of term babies	Similar
Stillbirth rate	Similar
Smoking status at time of delivery	Worse
AE attendances (0-4 years)	Worse
Percentage of 5 year olds with experience of visually obvious dental decay	Better

Indicator	Compared to England
Under 18s conception rate / 1,000	Similar
Emergency hospital admissions for asthma (< 19 yrs)	Better
Emergency hospital admissions for epilepsy (< 19 yrs)	Similar
Emergency hospital admissions for diabetes (< 19 yrs)	Worse
Hospital admissions for mental health conditions (0-17 years)	Better
Hospital admissions as a result of self-harm (10-24 years)	Similar
Hospital admissions due to substance misuse (15-24 years)	Similar
Hypertension: QOF prevalence (all ages)	Lower
Diabetes: QOF prevalence (17+)	Similar
CHD: QOF prevalence (all ages)	Lower
CKD: QOF prevalence (18+)	Lower
Stroke: QOF prevalence (all ages)	Lower
Deaths from circulatory disease, under 75 years	Similar
Deaths from all cancer, under 75 years	Similar
Cancer diagnosed at early stage (experimental statistics)	Not compared
Unplanned hospitalisation for chronic ACSC	Worse
Depression: Recorded prevalence (aged 18+)	Higher
Serious Mental Illness: QOF prevalence (all ages)	Lower

Indicator	Compared to England
Suicide rate (Persons)	Worse
Suicide rate (Male)	Worse
Estimated dementia diagnosis rate (aged 65 and over)	Not compared
Emergency hospital admissions due to falls (persons aged 65 and over)	Worse
Emergency hospital admissions for hip fracture (persons aged 65 and over)	Worse
Osteoporosis: QOF prevalence (50+)	Lower

SHEPPEY PCN

Indicator	Compared to England
Job Seekers Allowance rate (16-64 years)	Higher
Fuel poverty	Not compared
Life expectancy (Female)	Worse
Life expectancy (Male)	Similar
Estimated smoking prevalence (QOF)	Higher
Year 6: Prevalence of overweight (including obesity)	Worse
Admission episodes for alcohol-specific conditions	Better
Total number of prescribed antibiotic items per STAR-PU	Higher
Breast cancer screening coverage (females aged 50-70)	Better
Cervical cancer screening coverage (females aged 25-49)	Similar
Bowel cancer screening coverage (persons aged 60-74)	Similar
A&E attendances (0-4 years)	Worse
Emergency hospital admissions for asthma (< 19 yrs)	Worse
Hospital admissions as a result of self-harm (10-24 years)	Similar
CHD prevalence	Higher
Stroke prevalence	Similar
PAD prevalence	Similar
Heart failure prevalence	Higher

Indicator	Compared to England
AF prevalence	Similar
Hypertension prevalence	Higher
Cancer prevalence	Higher
Diabetes prevalence	Higher
COPD prevalence	Higher
Serious mental illness prevalence	Lower
Depression prevalence	Higher
Dementia prevalence	Similar
Premature mortality from all causes	Worse
Cancer mortality U75	Worse
CVD mortality U75	Worse
Osteoporosis prevalence 50+	Similar
Hip fracture adm 65+	Similar

SITTINGBOURNE PCN

Indicator	Compared to England
Job Seekers Allowance rate (16-64 years)	Similar
Fuel poverty	Not compared
Life expectancy (Female)	Similar
Life expectancy (Male)	Similar
Estimated smoking prevalence (QOF)	Higher
Year 6: Prevalence of overweight (including obesity)	Similar
Admission episodes for alcohol-specific conditions	Better
Total number of prescribed antibiotic items per STAR-PU	Higher
Breast cancer screening coverage (females aged 50-70)	Better
Cervical cancer screening coverage (females aged 25-49)	Better
Bowel cancer screening coverage (persons aged 60-74)	Better
A&E attendances (0-4 years)	Worse
Emergency hospital admissions for asthma (< 19 yrs)	Similar
Hospital admissions as a result of self-harm (10-24 years)	Similar
CHD prevalence	Lower
Stroke prevalence	Lower
PAD prevalence	Lower
Heart failure prevalence	Similar

Indicator	Compared to England
AF prevalence	Similar
Hypertension prevalence	Higher
Cancer prevalence	Higher
Diabetes prevalence	Similar
COPD prevalence	Similar
Serious mental illness prevalence	Lower
Depression prevalence	Higher
Dementia prevalence	Similar
Premature mortality from all causes	Similar
Cancer mortality U75	Similar
CVD mortality U75	Similar
Osteoporosis prevalence 50+	Similar
Hip fracture adm 65+	Better